

DUPIXENT[®]
(dupilumab) Injection 300mg

**HELP
HEAL
YOUR
SKIN
FROM
WITHIN™**

A WIN FOR SKIN IN

PRURIGO NODULARIS (PN)

**THE FIRST FDA-APPROVED
TREATMENT FOR ADULTS WITH PN**

Patient portrayal

INDICATION


DUPIXENT is a prescription medicine used to treat adults with prurigo nodularis (PN). It is not known if DUPIXENT is safe and effective in children with PN under 18 years of age.

Please see Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

PRURIGO NODULARIS STARTS BENEATH THE SKIN

Although the cause of Prurigo Nodularis is not entirely understood, PN is thought to begin inside the body where an abnormal relationship between the skin, nervous system, and immune system can lead to inflammation. This contributes to PN symptoms.

The intense cycle of itching and scratching and chronic inflammation leads to the development of nodules (bumps) and potentially lasting scars.



JENNIFER,
ACTUAL DUPIXENT PATIENT
Individual results may vary.

FACTS ABOUT PN



IT'S AN UNCOMMON DISEASE that is widely misunderstood and frequently misdiagnosed



ALMOST 311K ADULTS are impacted by PN in the USA



ABOUT 74% OF PATIENTS have an inadequate response to topical steroids



WOMEN MAY HAVE A HIGHER PREVALENCE OF PN

TALK TO A PN SPECIALIST IF YOU:

- ▶ HAVE PN NODULES THAT DO NOT GO AWAY
- ▶ NEED HELP BREAKING THE INTENSE CYCLE OF ITCHING
- ▶ HIDE YOUR SKIN FROM OTHERS



Scan the code or [click here](#) for a doctor discussion guide to review with your healthcare provider at your next appointment.

IMPORTANT SAFETY INFORMATION: Do not use if you are allergic to dupilumab or to any of the ingredients in DUPIXENT®.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.

LESS ITCHING AND FEWER BUMPS THAT'S A WIN FOR SKIN

DUPIXENT works under the skin, and may help bring balance between your skin, nervous system, and immune system by targeting a source of unwanted inflammation under your skin to help keep you one step ahead of PN symptoms. **DUPIXENT is the first FDA-approved treatment for adults with PN to help reduce itch and nodules.**



BROOKE,
ACTUAL DUPIXENT PATIENT
Individual results may vary.

"Prior to starting DUPIXENT, I was uncomfortable due to the many nodules on my arms and legs."

DUPIXENT:

- ▶ IS A BIOLOGIC THAT PROACTIVELY TREATS YOUR PN
- ▶ HELPS HEAL YOUR SKIN FROM WITHIN
- ▶ IS NOT A STEROID, CREAM, OR IMMUNOSUPPRESSANT
- ▶ **OVER 32,000*** ADULT PN PATIENTS ON THERAPY IN THE US

*IQVIA National Source of Business (NSOB) data as of June 2025.

IMPORTANT SAFETY INFORMATION: Before using DUPIXENT, tell your healthcare provider about all your medical conditions, including if you: have eye problems; have a parasitic (helminth) infection; are scheduled to receive any vaccinations. You should not receive a "live vaccine" right before and during treatment with DUPIXENT; are pregnant or plan to become pregnant. It is not known whether DUPIXENT will harm your unborn baby. A pregnancy registry for women who take DUPIXENT during pregnancy collects information about the health of you and your baby; are breastfeeding or plan to breastfeed. It is not known whether DUPIXENT passes into your breast milk.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).





RESULTS YOU CAN SEE AND FEEL THAT'S A WIN FOR SKIN



JENNIFER,
ACTUAL DUPIXENT PATIENT
Individual results may vary.

"My days were filled with a combination of physical discomfort, like itchiness and painful nodules."

PROVEN RESULTS IN ADULTS:

DUPIXENT was proven to reduce itch and help achieve clearer skin.

**IN TWO CLINICAL STUDIES WITH
311 ADULTS WITH PN:**

At 24 Weeks

**59% HAD NOTICEABLY
LESS ITCH**

59% on DUPIXENT vs 19% not on DUPIXENT

In as little as 12 weeks in one study

**37% EXPERIENCED
LESS ITCH**

37% on DUPIXENT vs 22% not on DUPIXENT

At 24 Weeks

**46% SAW SKIN CLEAR OR ALMOST
CLEAR OF PN NODULES**

46% on DUPIXENT vs 17% not on DUPIXENT

The most common side effects in patients with PN include: eye and eyelid inflammation, including redness, swelling, and itching, sometimes with blurred vision, herpes virus infections, common cold symptoms (nasopharyngitis), dizziness, muscle pain, diarrhea.

IMPORTANT SAFETY INFORMATION: Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).

ONE DOSE EVERY TWO WEEKS

DUPIXENT is taken as one dose every two weeks, after an initial loading dose of two injections. Here are other things to know:

- ▶ It's not taken every day
- ▶ It's a biologic medicine given under the skin (subcutaneous injection)
- ▶ Take it at home or at the doctor's office, after receiving training from your doctor.



PRE-FILLED PEN



PRE-FILLED SYRINGE

The recommended dosage for adult patients is an initial dose of 600 mg (two 300-mg injections) followed by 300 mg given every other week.


Use DUPIXENT exactly as prescribed by your healthcare provider. Your healthcare provider will tell you how much DUPIXENT to inject and how often. **Do not** try to prepare and inject DUPIXENT until you or your caregiver have been trained by your healthcare provider.



Scan the code or [click here](#) to learn more about how to inject DUPIXENT.

IMPORTANT SAFETY INFORMATION: Especially tell your healthcare provider if you are taking oral, topical, or inhaled corticosteroid medicines or if you have PN and asthma and use an asthma medicine. **Do not** change or stop your other medicines, including corticosteroid medicine or other asthma medicine, without talking to your healthcare provider. This may cause other symptoms that were controlled by those medicines to come back.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).



ROGER,
ACTUAL DUPIXENT PATIENT
Individual results may vary.

HELPFUL RESOURCES FOR YOU, EVERY STEP OF THE WAY

DUPIXENT MyWay is a patient support program designed to help you start DUPIXENT as quickly as possible and stay on track with your treatment.

ENROLLING IN **DUPIXENT MYWAY**[®] IS FREE AND SIMPLE

Contact us directly at 1-844-DUPIXENT
(1-844-387-4936) Option 1, Monday–Friday,
8 AM–9 PM ET.

Once enrolled in the program, you'll have a dedicated Case Manager who can provide:



Help with understanding your insurance coverage for DUPIXENT



Financial assistance, including copay support, if eligible



Connection to a **DUPIXENT MyWay** Nurse Educator who can offer supplemental injection training in person, virtually, or by the phone



Tips and tools to help you manage treatment

IMPORTANT SAFETY INFORMATION: DUPIXENT can cause serious side effects, including: Allergic reactions. DUPIXENT can cause allergic reactions, including skin reactions, that can sometimes be severe. Stop using DUPIXENT and tell your healthcare provider or get emergency help right away if you get any of the following signs or symptoms: breathing problems or wheezing, swelling of the face, lips, mouth, tongue, or throat, fainting, dizziness, feeling lightheaded, fast pulse, fever, hives, skin rash, including rash that looks like a bullseye or painful red or blue bumps under the skin, general ill feeling, itching, swollen lymph nodes, nausea or vomiting, joint pain, or cramps in your stomach area.

OPTIONS FOR FINANCIAL SUPPORT



The **DUPIXENT MyWay Copay Card Program** may help eligible patients cover out-of-pocket costs for DUPIXENT.*

The **DUPIXENT MyWay Patient Assistance Program**[†] may be able to help you if you do not have health insurance, are experiencing difficulty paying for your DUPIXENT treatment, or have Medicare Part D.

Find out if you qualify by speaking with a **DUPIXENT MyWay** Case Manager at 1-844-DUPIXENT (1-844-387-4936) Option 1.



Scan the code or [click here](#) to learn more about how to inject DUPIXENT.

*Terms and conditions apply.

[†]Eligible commercial patients and Medicare patients without the Part D (pharmacy) benefit may receive free shipments of up to an 84-day supply of DUPIXENT (for up to 12 months). Patients may reapply after 12 months if they still meet the eligibility criteria. Patients with Medicare Part D who meet eligibility criteria must reapply for the Patient Assistance Program each calendar year, or by December 31st for eligibility consideration for the following year. The assistance period for Medicaid patients varies based on the eligibility criteria. Please note that DUPIXENT MyWay reserves the right to make eligibility determinations, monitor participation, ensure equitable product availability, and modify or discontinue the DUPIXENT MyWay Patient Assistance Program at any time without notice.

IMPORTANT SAFETY INFORMATION: DUPIXENT can cause serious side effects, including (cont'd): **Eye problems.** Tell your healthcare provider right away if you have any new or worsening eye problems, including eye pain or changes in vision, such as blurred vision. Your healthcare provider may send you to an ophthalmologist for an eye exam if needed.

Joint aches and pain. Some people who use DUPIXENT have had trouble walking or moving due to their joint symptoms, and in some cases needed to be hospitalized. Tell your healthcare provider about any new or worsening joint symptoms. Your healthcare provider may stop DUPIXENT if you develop joint symptoms.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information including Patient Information.



DARICHA,
ACTUAL DUPIXENT PATIENT
Individual results may vary.

A WIN FOR SKIN IN PRURIGO NODULARIS (PN)

ASK YOUR DERMATOLOGIST ABOUT DUPIXENT.

For more information visit [DUPIXENT.com](https://www.dupilumab.com) or call
1-844-DUPIXENT (1-844-387-4936).

IMPORTANT SAFETY INFORMATION:

The most common side effects in patients with prurigo nodularis include eye problems, including eye and eyelid inflammation, redness, swelling, itching, and blurred vision; herpes virus infections; common cold symptoms (nasopharyngitis); dizziness; muscle pain; and diarrhea.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of DUPIXENT. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Use DUPIXENT exactly as prescribed by your healthcare provider. It's an injection given under the skin (subcutaneous injection). Your healthcare provider will decide if you or your caregiver can inject DUPIXENT. Do not try to prepare and inject DUPIXENT until you or your caregiver have been trained by your healthcare provider.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#) including [Patient Information](#).

sanofi | **REGENERON**[®]

© 2026 Sanofi and Regeneron Pharmaceuticals, Inc. All rights reserved. DUPIXENT and Dupixent MyWay are registered trademarks of Sanofi or an affiliate.

US.DUP.26.04.0297 05/26



SCAN THE CODE OR [CLICK HERE](#)
TO LEARN MORE ABOUT HOW
DUPIXENT CAN HELP